Dissolution Pathways and Mother-Child Relationship Quality Among Adolescents

Elizabeth S. Howe-Huist (eshoweh@bgsu.edu)

Department of Sociology Bowling Green State University



Background

- Children in cohabiting and married stepfamilies experience higher rates of instability, yet there has been scarce research on outcomes associated with these types of dissolutions
- Parental divorce is consistently associated with worse outcomes for children

Research Questions

- How does mother-child relationship quality change from pre-dissolution to post-dissolution, and do these changes vary by dissolution pathway?
- Does the loss of economic and parental resources explain the change in mother-child relationship quality across different dissolution pathways?


Data and Method

- National Longitudinal Study of Adolescent to Adult Health (Add Health) Waves I & II
- Analytic Sample: n= 902
 - Includes those who reside with their mother and either their father, mother's husband, or mother's partner in Wave I, and indicated a change to not having the same male figure in the home by Wave II.
- Ordinary Least Squares Regression

Dissolution Pathways

- Dissolution pathway: A dissolution or divorce a child experiences between their parental figures
 - Divorce of biological parents (Biological Divorce, n= 526)
 - Divorce of biological mother and stepfather (Stepparent Divorce, n= 201)
 - Dissolution of biological mother and cohabiting partner (Cohabitation Dissolution, n= 175)


Figure 1. Sample by Dissolution Pathway




Mother-Child Relationship Quality

- Mother-child relationship quality is assessed using five questions asked of adolescents in Wave I and Wave II (α =.83 WI & α =.85 WII)
- Change in mother-child relationship quality (Wave II-Wave I)

Figure 2. Changes in Mother-Child Relationship Quality



Logistic Regression



Conclusion and Future Research

- The change in mother-child relationship quality does not significantly differ across the three dissolution pathways
- Examine other areas of well-being including academic outcomes, delinquent behaviors, and psychological well-being
- Explore long-term well-being of adolescents who have experienced different dissolution pathways

This research was supported in part by the Center for Family and Demographic Research, Bowling Green State University, which has core funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (P2CHD050959). This research uses data from Add Health, a program project directed by Kathleen Mullan Harris and designed by J. Richard Udry, Peter S. Bearman, and Kathleen Mullan Harris at the University of North Carolina at Chapel Hill, and funded by grant P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 23 other federal agencies and foundations. Special acknowledgment is due Ronald R. Rindfuss and Barbara Entwisle for assistance in the original design. Information on how to obtain the Add Health data files is available on the Add Health website http://www.cpc.unc.edu/addhealth. No direct support was received from grant P01-HD31921 for this analysis.